


ferti•lome®



F-Stop™

Specialty Fungicide

A Granular Product Containing a Systemic Protectant and Curative Fungicide for the Control of Diseases in Established Home Lawns and in Ornamental Turfgrass

- Cures and Prevents All Major Lawn Diseases
- Prevents Over 15 Major Lawn Diseases
- For Use On All Types Of Home Lawns
- One Application Protects For Up To 4 Weeks

ACTIVE INGREDIENT:	
myclobutanil: a-butyl-a-(4-chlorophenyl)-1-H-1,2,4 triazole-propanenitrile	0.39%
OTHER INGREDIENTS:	99.61%
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

NET WEIGHT 20 LBS. (9.07 KG)

ferti•lome® F-Stop™

ferti•lome® F-Stop™

10768

ferti•lome®

F-Stop™



ferti•lome® F-Stop™

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes Moderate Eye Irritation.

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Not for use on turfgrass being grown for sale or other commercial use as sod, or for commercial seed productions, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-992-5994 for disposal instructions.

Never place unused product down any indoor or outdoor drain.

Notice: Read the entire label. Use only according to label directions. Before using this product, read "Warranty Disclaimer," "Inherent Risks of Use," and "Limitation of Remedies" at end of Directions for Use. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call **1-800-992-5994**

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

HOW THIS PRODUCT WORKS

ferti•lome® F-STOP™ is a systemic, protectant and curative fungicide for the control of listed diseases in established home lawns and ornamental turfgrass. Optimum disease control is achieved when this product is applied to established turfgrass in a regularly scheduled preventative program. Use this product in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Use diagnostic kits, extension experts, or other identification methods when developing disease control strategies.

HOW TO APPLY

Apply **ferti•lome® F-STOP™** uniformly over the turfgrass area using a properly calibrated drop or rotary-type spreader designed to apply granules. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Check frequently to make sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps during application. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands.

Wash hands with soap and water promptly after use.

Do not allow people or pets to contact treated area until after product dust has settled into the turfgrass, or if watered in, after the turfgrass surface in the treated area has dried.

Disease	ferti•lome® F-STOP™ (lb/1,000 sq ft)	Application Interval/Timing (Days)	Directions	Restrictions
Anthracnose Red Thread Septoria Leaf Spot	4	14 - 21	Apply when conditions are favorable for disease development.	Do not apply more than 46 pounds of product per 1,000 sq. ft. per year. For Nassau and Suffolk counties in New York State, do not apply more than 11.5 lb of this product per 1,000 sq ft per year.
Brown Patch		14	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent	
Copper Spot Zonate Leaf Spot			Apply when conditions are favorable for disease development.	
Crown Rot Leaf Spot Melting-Out			Apply when conditions are favorable for disease development. For crown rot, water in with 3 to 4 gallons of water per 1,000 sq. ft. to increase penetration to crown and roots.	
Dollar Spot		14 - 28	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for Dollar Spot control before rotating to a registered fungicide with a different mode of action	
Fusarium Blight		14 - 21	Apply when conditions are favorable for disease development.	
Fusarium Patch (Pink Snow Mold)		21 - 28	Apply when conditions are favorable for disease development.	
Gray Leafspot		14	Apply prior to snow cover.	
Leaf Smuts			Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.	
Necrotic Ring Spot		Spring: 28	Make applications on a preventative basis in early to mid-spring.	
Necrotic Ring Spot		Fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant.	
Powdery Mildew Rusts		14 - 28	Apply when conditions are favorable for disease development.	
Summer Patch			Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local Turfgrass Extension Experts. Water in with at least 3 to 4 gallons of water per 1,000 sq. ft. to increase penetration to crown and roots.	

WHEN TO APPLY

Minimize the interval between applications of this product when conditions are favorable for disease development. Unless otherwise specified, when disease pressure is high or when used as a curative, use higher rates of **ferti•lome® F-STOP™** and shorter application intervals. Under light to moderate disease pressure apply this product at the low use rate and/or longer application intervals. To avoid pick-up, lightly irrigate treated areas soon after application. On short cut bentgrass (1/2 inch or less) when temperatures are above 80°F, apply only to dry foliage.

HOW MUCH TO APPLY

Optimum disease control is achieved when **ferti•lome® F-STOP™** is applied in a preventative disease control program at a rate of 4 lb per 1,000 sq. ft. See the following table for specific application rates for various diseases. Under any circumstances, do not apply more than 46 lb of this product per 1,000 sq. ft. per year.

SPREADER SETTINGS AS A GUIDE FOR CALIBRATION

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)
Scotts®/Republic Accugreen (Drop)	4 1/4
Scotts®/Republic Speedy Green (Broadcast)	3 3/4
ferti•lome® /EarthWay Ev-N-Spred (Broadcast)	14

TERMS AND CONDITION OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the Seller for a full refund of purchase price paid. Otherwise, use by the Buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use, and Limitations of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description of the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the Inherent Risks set forth below. Seller **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by the Buyer.

LIMITATION OF REMEDIES

To the extent permitted with law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- Refund of purchase price paid by Buyer or user for product bought, or
- Replacement of amount of product used.

Seller shall not be liable for the losses of damages resulting from the handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use above and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™&® Trademarks of Corteva Agriscience and its affiliated companies

10768 Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No. 62719-461-7401
EPA Est. No. 7401-TX-01
Product Questions? 855-270-4776

ferti•lome® F-Stop™

6D10768-2

